

The Dimensionality of the Geiger Nuttall Law: a simple but interesting note

Pavle I. Premović

Laboratory for Geochemistry, Cosmochemistry and Astrochemistry

e-mail: pavleipremovic@yahoo.com

Abstract. It appears that the natural logarithm form of the empirical Geiger-Nuttall law is not dimensionally correct. It is suggested a new type of this form as well four new non-exponential forms of this law.

Keywords: Geiger-Nuttall law, dimensionality, half-life, alpha decay, alpha emitter.

One of the milestones of quantum mechanics was the formulation of the Geiger-Nuttall law [1, 2]. This law states that there is a simple empirical relation between the α -decay energy Q_α and the half-life of heavy alpha (α)-emitter $t_{1/2}$

$$\log t_{1/2} = A + B/\sqrt{Q_\alpha}$$

where A and B are constants. Let us convert this equation into a natural logarithm (LN) form

$$\ln t_{1/2} = A + B/\sqrt{Q_\alpha}$$

where $a = 2.303A$ and $b = 2.303B$.¹ The kinetic energy, E_α , of the emitted α -particle is slightly less (about 0.4 % for heavy α -emitters, such as the uranium α -emitters) than Q_α . Therefore, the last equation can be written as follows

$$\ln t_{1/2} = a + b/\sqrt{E_\alpha} \quad \dots (1).$$

After the conversion of this equation into the exponential form we have

$$t_{1/2} = e^{a + b/\sqrt{E_\alpha}} \quad \dots (2).$$

Obviously, the dimension² of the left side of this expression is T but its right side is dimensionless. Hence, eqn. (2) is dimensionally incorrect. The simplest way to make this equation dimensionally correct is to multiply its right side with the constant θ equals or close to 1 and expressed in time unit of $t_{1/2}$ (e. g. seconds)

$$t_{1/2} = e^{a + b/\sqrt{E_\alpha}} \times \theta$$

¹ $\ln t_{1/2} = 2.303 \log t_{1/2}$

² Just to remind the reader that M, L and T are the symbols for basic dimensions: mass, length and time, respectfully.

Since $\ln\theta$ is equal or close to 0 the LN form of this equation is identical to the initial eqn. (1) of the Geiger-Nuttall law.

There is another way to resolve the above dimensionality issue. Instead the eqn. (1) derived from empirical Geiger-Nuttall law we may propose the following equation

$$t_{1/2} = ab/\sqrt{E_\alpha} \quad \dots (4).$$

The LN form of this equation is $\ln t_{1/2} = \ln a + \ln(b/\sqrt{E_\alpha})$ where a and b are the constants. The last equation is a linear form of $t_{1/2}$ vs. $1/\sqrt{E_\alpha}$. After some rearrangement of eqn. (4) we get

$$t_{1/2} = a\sqrt{(b^2/E_\alpha)}.$$

Keeping in the mind that the dimension of E_α^3 is ML^2T^{-2} , a dimensional analysis shows that this equation is dimensionally correct for the fourth following cases: 1) if the dimension of a is L then the dimension of b^2 must be M; 2) if the dimension of a is T then the dimension of b^2 must be ML^2T^{-2} , i. e. b^2/E_α must be dimensionless; 3) if the dimension of a is M then the dimension of b^2 must be $M^{-1}L^2$; and finally 4) if a is dimensionless then the dimension of b^2 must be ML^2 .

The detailed analysis of each of the above cases, including eqns. (1) and (2), will be subject of author's further study.

Recently, Premović [3] derived a new mathematical form for the Geiger-Nuttall law. This is expressed by the following equation

$$t_{1/2} = u \times w \times 1/\sqrt{E_\alpha}$$

where u and w are the constants. According to Premović [7], the dimension of u is L (analogous the above case 1) and the dimension of $(w \times 1/\sqrt{E_\alpha})$ is $[L^{-1} \times T]$ or the dimension of w is M. Therefore, the left side of this expression is dimensionally consistent with its right side.

References

- [1] H. Geiger and J.M. Nuttall, *The ranges of the α particles from various radioactive substances and a relation between range and period of transformation*. Phil. Mag., 22, 613-621 (1911).
- [2] H. Geiger and J.M. Nuttall, *The ranges of α particles from uranium*, Phil. Mag., 23, 439-445 (1912).
- [3] P. I. Premović, *The Lifetimes of ^{238}U Nuclei, the Escaping Attempts of Their Alpha Particles and the Geiger-Nuttall Law*. The General Science Journal, June 2019.

³ Since $E_\alpha = 1/2mv^2$ (where m is the mass of the α -particle and v is its speed) and the dimension of v is LT^{-1} then the dimension of E_α is ML^2T^{-2}