

## The Big Bang Universe and the Principle of Energy Conservation

Pavle I. Premović,  
Laboratory for Geochemistry, Cosmochemistry&Astrochemistry,  
University of Niš, 18000 Niš, Serbia

### Abstract

The energy of light measured by an observer comoving with a nearby or distant galaxy emitting this light would be  $N$  (a rational number) times larger than the energy of galaxy's light measured by Earth's observer. The difference between these two energies in the case of nearby galaxies would be quantized by photon quantum energy  $\varepsilon = hH_0$  where  $h$  is Planck's constant and  $H_0$  is Hubble's constant.

**Keywords:** Big Bang, Universe, redshift, Doppler effect, energy.

### Introduction

The Big Bang theory states that the observable Universe began 13.8 Gy ago, in an enormous and extremely fast expansion. General relativity claims that this expansion is an expansion of space (or better space-time) itself and still is (uniformly) occurring. According to this theory the wavelength of light coming from nearby or distant galaxies<sup>1</sup> increases or shows cosmological redshift (or simply redshift).

Let us denote with  $\lambda$  (or  $\nu$ ) the wavelength (or frequency) of light emitted by the (nearby or distant) galaxy source (or generated by the same source on the Earth) and  $\lambda_G$  (or  $\nu_G$ ) the wavelength (or frequency) of light or measured by an Earth observer. As we noted above, if  $\lambda_G > \lambda$  (or  $\nu > \nu_G$ ) then galaxy's light is redshifted.

The redshift of a galaxy  $z_G$  in cosmology is characterized by the relative difference between the observed wavelength  $\lambda_G$  and emitted wavelength  $\lambda$  of light (or in general electromagnetic radiation) sourced by nearby or distant galaxies. This is mathematically expressed by the following equation

$$z_G = (\lambda_G - \lambda)/\lambda \quad \dots (1).$$

---

<sup>1</sup> We define nearby galaxies as those galaxies whose redshift  $z_G$  is from 0.001 to 0.1 (or  $0.001 \leq z_G \leq 0.1$ ) and distant galaxies those having  $z_G > 0.1$  [1, 2]. Of course, there is no sharp boundary between nearby and distant galaxies.

The vast majority of nearby and all distant galaxies<sup>2</sup> show redshift in their spectra.

In contrast to the Big Bang model, the ultraviolet surface brightness data of galaxies, over a very wide redshift range imply that the observable Universe is the non-expanding (Euclidean) Universe (NEEU) [3, and references therein]. A detailed analysis of the gamma-ray burst sources performed by Sanejouand [4] supports this view suggesting that this Universe has been Euclidean and static over the last 12 Gy. To explain redshift, Premović [1] hypothesized that the speed of light emitted by nearby/distant galaxies in NEEU is superluminal.

The conservation of energy (the Principle energy of conservation) is one of the fundamental laws of physics which is not violated by any known process. For example, in quantum physics, energy is conserved. In contrast, one of the problems which the Big Bang Universe is facing is the apparent violation of this law by the cosmological expansion.

Another problem which has received little attention despite its importance is the fact that we do not know the recessional speed of any nearby or distant galaxies at all (for example, see <https://www.loop-doctor.nl/hubble-and-humason-measured-redshift/>). Moreover, we do not know the distance between the Earth and these galaxies except for a few nearby so-called “megamaser” galaxies (1). In further text, we will assume that a peculiar speed of nearby/distant galaxies is negligible. Indeed, almost all these galaxies have such peculiar speeds [5].

## Derivation and Discussion

Let us assume that a nearby or distant galaxy is moving from the Earth emitting the photon toward Earth’s observer and who performs a measurement on it. Assume that the frequency of this photon is  $\nu_G$  and its energy, according to Planck’s equation, is  $E_G = h\nu_G$  where  $h$  is Planck’s constant ( $6.63 \times 10^{-34}$  J sec). This is only the energy that we could assign to the photon in question following the conservation energy law. On the other hand, an observer comoving with a nearby or distant galaxy would measure a higher frequency  $\nu$  and calculate higher energy,  $E = h\nu$ , which is also conserved. In other words, the energy of the photon emitted by these galaxies in the Big Bang Universe is conserved in the two different (galaxy and Earth) reference frames. This is in accord with Special relativity which allows that the observers in different frames of reference can measure different energies for the same event.

Denote with  $\mathcal{N}_G$  the number of periods  $T_G (= 1/\nu_G)$  for the light emitted by a nearby or distant galaxy to the Earth. Analogously, denote by  $\mathcal{N}$  the number periods  $T (= 1/\nu)$  of this light but viewed by an observer comoving with any of these galaxies. The distance,  $D_G$ , between the Earth and a nearby or distant galaxy, is identical regardless of whether the Earth or galaxy is moving away. So, the number of periods is identical. Mathematically speaking we state

$$D_G/c = \mathcal{N}/\nu = \mathcal{N}_G/\nu_G \quad \dots (2).$$

---

<sup>2</sup> Almost all nearby and all distant galaxies are redshifted; they are moving away from the Earth. There are about 100 galaxies that have blueshifts and are heading towards the Earth. Most of these are Local Group (dwarf) galaxies.

where  $c$  ( $= 2.99792 \times 10^8$  m sec<sup>-1</sup>) is the speed of light. Of course,  $\mathcal{N}_G$  and  $\mathcal{N}$  are extremely large natural numbers.<sup>3</sup> After some rearrangement of this equation, we get

$$\mathcal{N}v_G = \mathcal{N}_G v. \quad \dots (3).$$

Since  $v > v_G$ , then  $\mathcal{N} > \mathcal{N}_G$ .

Elementary physics shows that

$$v = c/\lambda \text{ and } v_G = c/\lambda_G$$

Combining these two formulas with eqn. (2) we have

$$D_G = \mathcal{N}\lambda = \mathcal{N}_G\lambda_G$$

Since  $\mathcal{N}$  and  $\mathcal{N}_G$  are extremely large natural numbers then the ratio  $(\mathcal{N}/\mathcal{N}_G)$  is a rational number that will be denoted as  $N$  or  $N = \mathcal{N}/\mathcal{N}_G$ . After a bit of algebra we

$$\lambda_G/\lambda = \mathcal{N}/\mathcal{N}_G$$

or

$$\lambda_G/\lambda = N \quad \dots (4)$$

Multiplying eqn. (3) with  $h$  and after a bit of algebra we get

$$hv/hv_G = E/E_G = \mathcal{N}/\mathcal{N}_G.$$

For convenience, we write

$$E = (\mathcal{N}/\mathcal{N}_G)E_G$$

or

$$E = NE_G \quad \dots (5).$$

Eqn. (2) and all subsequently derived equations are valid for all nearby/distant galaxies with the identical source of emitting light.

---

<sup>3</sup> Each large rational number (say  $\geq 10000$ ) can be approximated with a corresponding natural number since the approximation error is small ( $\leq 0.05\%$ ). In the case of nearby or distant galaxies, we are dealing with an extremely large rational number of the periods, therefore the approximation error is extremely small – i. e. completely negligible.

In summary, the energy (or frequency) of light emitted by a nearby or distant galaxy measured by an observer commoving with the galaxy would be  $N$  times larger than the energy (or frequency) of this light measured by an Earth observer. In contrast, the wavelength of this light measured by an Earth observer would be (apparently)  $N$  times its wavelength determined by an observer commoving with the galaxy.

Let us denote with  $\Delta E$  a difference of energy between  $E = hv$  and  $E_G = hv_G$  then

$$\Delta E = E - E_G = hv - hv_G \quad \dots (6).$$

If we combine the middle part of this equation with eqn. (5) we get

$$\Delta E = (N - 1)E_G.$$

Cosmologists consider Hubble's law as a direct consequence of the expansion of Universe from the initial Bing Bang. This law states that there is a linear relationship between the distance  $D_G$  to nearby galaxies and the redshift  $z_G$  of their light. It can be expressed as

$$z_G = D_G H_0 / c \quad \dots (7)$$

where  $H_0$  is Hubble's constant representing the constant rate of the Universe expansion and it ranges from  $50 \text{ km sec}^{-1} (\text{Mpc})^{-1}$ -  $100 \text{ km sec}^{-1} (\text{Mpc})^{-1}$ . We usually use  $H_0 = 72 \text{ km sec}^{-1} (\text{Mpc})^{-1}$ .

Combining eqn. (1) and eqn. (7) we arrive at

$$\Delta E = (D_G / \lambda_G) h H_0 = \mathcal{N}_G h H_0 \quad \dots (8).$$

Combining this equation and eqn. (7) we simply derive

$$\Delta E \lambda_G / z_G = hc$$

As we noted above,  $\mathcal{N}_G$  is an extremely large natural number so the difference of energy  $\Delta E$  is quantized. This possibility was just mentioned in our previous communication [6]. Then we defined  $hH_0$  as the photon quantum of energy  $\varepsilon$  or

$$\varepsilon = hH_0.$$

Using the above value for  $H_0$  we estimated that  $\varepsilon = 1.5 \times 10^{-51} \text{ J}$ . For further details see [6].

All equations, derived using eqn. (7), are only valid for nearby galaxies.

It is here worth noting that the cosmological redshift of nearby/distant galaxies is given by

$$z_G + 1 = a(t) / a(t_G)$$

where  $a(t)$  is the scale factor at the cosmic time of emission of light by a nearby or a distant galaxy and  $a(t_G)$  is the scale factor later at cosmic time  $t_G$  of Earth's observer. A further consideration is outside the scope of this communication.

## References

- [1] P. I. Premović, *Distant galaxies in the non-expanding (Euclidean) Universe: the light speed redshift*. The General Science Journal, May 2020.
- [2] ] P. I. Premović, *Nearby and distant galaxies: a brief note*. The General Science Journal, October 2020.
- [3] E. J. Lerner, *Observations contradict galaxy size and surface brightness predictions that are based on the expanding universe hypothesis*. Monthly notices the Royal Astron. Soc. (MNRAS) 477, 3185-3196 (2018).
- [4] Y. –H Sanejouand, *About some possible empirical evidences in favor of a cosmological time variation of the speed of light*. Europhys. Lett., 88, 59002 (2009).
- [5] T. M. Davis and M. I. Scrimgeour, *Deriving accurate peculiar velocities (even at high redshift)*. MNRAS, 442, 1117–1122 (2014).
- [6] P. I. Premović, *The photon quantum of energy of the observable universe*. The General Science Journal, October 2020.