

## New prime $K$ -tuple theorem (3)

$$P, jP + j + 1 (j = 1, \dots, k)$$

Chun-Xuan Jiang

P. O. Box 3924, Beijing 100854, P. R. China

[Jiangchunxuan@vip.sohu.com](mailto:Jiangchunxuan@vip.sohu.com)

### Abstract

Using the Jiang function, we prove that for every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $jP + j + 1$  is prime.

### Theorem

$$P, jP + j + 1 (j = 1, \dots, k). \quad (1)$$

For every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $jP + j + 1$  is prime.

**Proof.** We have Jiang function [1, 2]

$$J_2(\omega) = \prod_p (P - 1 - \chi(P)), \quad (2)$$

where  $\omega = \prod_p P$ ,

$\chi(P)$  is the number of solutions of congruence

$$\prod_{j=1}^k (jq + j + 1) \equiv 0 \pmod{P}, \quad (3)$$

where  $q = 1, \dots, P - 1$ .

From (3) we have

If  $P \leq k + 1$  then  $\chi(P) = P - 2$ , if  $k + 1 < P$  then  $\chi(P) = k$ .

From (3) and (2) we have

$$J_2(\omega) = \prod_{k+1 < P} (P - k - 1) \neq 0. \quad (4)$$

We prove that for every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $jP + j + 1$  is prime.

We have the best asymptotic formula [1, 2]

$$\pi_{k+1}(N, 2) = \left| \{ P \leq N : jP + j + 1 = \text{prime} \} \right| \sim \frac{J_2(\omega) \omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N}. \quad (5)$$

The author took a day to write this paper.

### References

- [1] Chun-Xuan Jiang, Jiang's function  $J_{n+1}(\omega)$  in prime distribution.  
<http://www.wbabin.net/math/xuan2.pdf> (<http://vixra.org/pdf/0812.0004v2.pdf>)
- [2] Chun-xuan Jiang, The Hardy-Littlewood prime  $k$ -tuple conjecture is false.  
[http:// www. wbabin.net/math/xuan77.pdf](http://www.wbabin.net/math/xuan77.pdf)